

A PKU TREATMENT THAT HELPS FIND

LESS LIMITS. MORE POSSIBILITIES.

See what
SEPHIENCE
can offer.



IMPORTANT SAFETY INFORMATION

What is SEPHIENCE?

SEPHIENCE™ (sepiapterin) is a prescription medicine used to lower blood levels of phenylalanine (Phe) in adults and children 1 month of age and older with a certain type of Phenylketonuria (PKU). SEPHIENCE is used along with a Phe-restricted diet. It is not known if SEPHIENCE is safe and effective in children younger than 1 month of age.

What is the most important information I should know about SEPHIENCE?

SEPHIENCE may cause serious side effects, including:

- **increased bleeding problems.** SEPHIENCE may increase your risk of bleeding. Tell your healthcare provider if you develop any symptoms of bleeding including:
 - bruising or red or purple skin marks
 - menstrual bleeding that is heavier than normal
 - bleeding longer than usual or that does not stop
- **low blood Phe (hypophenylalaninemia).** Low blood Phe levels are common during treatment with SEPHIENCE and can be severe.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

Navigating life with PKU

Living with Phenylketonuria—better known as PKU—comes with daily challenges, like managing phenylalanine (Phe) levels. SEPHIENCE is used to help manage Phe levels in people with PKU, as part of a Phe-restricted diet, allowing you to focus on what's important to you.

What is PKU?

PKU is a genetic disorder characterized by the body's inability to break down an amino acid called Phe. When we consume high-protein foods, the body breaks it down into smaller components like amino acids. In people with PKU, the body has trouble processing Phe properly, causing it to build up in the blood—which can lead to irreversible brain damage and other serious health problems.

What's it called when Phe is high?

It's called hyperphenylalaninemia—when too much Phe builds up in the blood.



What's the recommended Phe range?

Experts recommend keeping Phe $\leq 360 \mu\text{mol/L}$ ($\leq 6 \text{ mg/dL}$) for lifelong brain health.



Diet matters. People with PKU need to avoid high-protein foods like meat, fish, eggs, dairy products, and nuts, as these can raise Phe levels.

IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)

The most common side effects of SEPHIENCE include: diarrhea, headache, stomach (abdominal) pain, yellow or orange stool (feces), and throat pain.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of SEPHIENCE. Call your healthcare provider for medical advice about side effects.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

What is SEPHIENCE™?

SEPHIENCE is a prescription medicine used along with a Phe-restricted diet to lower blood levels of Phe in adults and children 1 month of age and older with PKU. It contains sepiapterin, which your body converts into tetrahydrobiopterin (BH_4)—a key molecule that supports the enzyme that breaks down Phe and reduces its buildup.

How SEPHIENCE works




1 Taken as an oral powder



2 Converts to natural BH_4 , which aids Phe breakdown



3 Reduces Phe levels to help manage PKU



SEPHIENCE doesn't just add BH_4 —it helps your body make its own, supporting your natural process.



IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)

Before taking or giving SEPHIENCE, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if SEPHIENCE will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if SEPHIENCE passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take SEPHIENCE.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

Results with SEPHIENCE

SEPHIENCE was studied in APHENITY, a two-part clinical trial of people aged 1-61 years old living with PKU. During Part 1 of the trial, 156 participants received their age- and weight-specific dosage. In Part 2 of the trial, a total of 98 people with PKU, aged 2 years and older, who responded to SEPHIENCE in Part 1 were randomly assigned to receive either SEPHIENCE or a placebo for 6 weeks.

In Part 1 of the clinical trial,

SEPHIENCE began lowering blood Phe levels for many participants in the clinical trial after just **2 weeks**.

66%

had $\geq 30\%$ reduction in Phe levels

73% had $\geq 15\%$ reduction in Phe levels



IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. SEPHIENCE and other medicines may affect each other causing side effects. SEPHIENCE may affect the way other medicines work, and other medicines may affect how SEPHIENCE works. Especially tell your healthcare provider if you take:

- a medicine that contains levodopa
- an antifolate medicine
- another medicine also used to lower Phe
- medicines used to treat erectile dysfunction (phosphodiesterase 5 inhibitors or PDE5)

Please see full Important Safety Information on [pages 12-13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

In Part 2 of the clinical trial,

SEPHIENCE™ led to a significant reduction in blood Phe levels compared to placebo after **6 weeks**.

63%

reduction in Phe levels in the study population

Individual results may vary.



IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)

Know the medicines you take. Keep a list of them with you to show your healthcare provider and pharmacist when you get a new medicine.

How should I take or give SEPHIENCE?

- Take or give SEPHIENCE exactly as your healthcare provider tells you. Your healthcare provider will tell you how much SEPHIENCE to take or give.
- Your healthcare provider may change your or your child's dose of SEPHIENCE depending on your weight and age.
- Take or give SEPHIENCE 1 time each day with food. You must take or give more food after the SEPHIENCE dose.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

Additional results with SEPHIENCE

Results across different types of PKU


As part of the APHENITY study, SEPHIENCE was studied in people with different types of PKU, including those with classical PKU and those who had not responded to BH_4 treatment in the past. Classical PKU is the most severe form of PKU. People with classical PKU have had very high levels of blood Phe ($\geq 1200 \mu\text{mol/L}$) at some point in their lives, even if their levels are lower now. In both groups, SEPHIENCE significantly lowered the amount of Phe in the blood, showing it worked across a wide range of people with PKU.

For people with classical PKU

Blood Phe was reduced by

69%

on SEPHIENCE



For people not helped by BH_4 treatment in the past

Blood Phe was reduced by

54%

on SEPHIENCE

These findings help demonstrate the potential for SEPHIENCE to support people with PKU regardless of prior treatment experience or PKU type.

IMPORTANT SAFETY INFORMATION (cont'd)

How should I take or give SEPHIENCE? (cont'd)

- SEPHIENCE comes as a packet containing oral powder.
 - For SEPHIENCE doses **less than** 1,000 mg: mix SEPHIENCE oral powder with water or apple juice before giving or taking.
 - For SEPHIENCE doses 1,000 mg or **greater**: SEPHIENCE oral powder can be mixed with water, apple juice, strawberry jam, or applesauce before taking or giving.
 - **See the detailed Instructions for Use that comes with SEPHIENCE for information about the correct way to prepare, take, or give a dose of SEPHIENCE oral powder.**

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

Achieving recommended Phe levels

SEPHIENCE™ was studied to see how well it helps people with PKU reach blood Phe level targets recommended by experts. These targets, set by the American College of Medical Genetics and Genomics, recommend maintaining a blood Phe level of ≤ 360 $\mu\text{mol/L}$ (or ≤ 6 mg/dL).

In the APHENITY trial:

84%

of participants taking SEPHIENCE reached that expert-recommended target

9% of participants who took placebo achieved the same goal

Maintaining Phe levels within this recommended range is important for supporting brain development and long-term cognitive health.

Individual results may vary. This information is based on clinical trial data but is not included in the approved full Prescribing Information for SEPHIENCE. Talk to your healthcare provider if you have any questions.

IMPORTANT SAFETY INFORMATION (cont'd)

How should I take or give SEPHIENCE? (cont'd)

- If you or your child miss a dose of SEPHIENCE, take or give it as soon as possible. Do not take 2 doses in a day. Continue with your normal dosing schedule the following day.
- During treatment with SEPHIENCE, you or your child will have regular check-ups with your healthcare provider.
- Your healthcare provider will do blood tests before and during your treatment with SEPHIENCE to check your blood Phe levels.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).



Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

Additional results with SEPHIENCE

APHENITY Extension Study: Effects on diet in PKU

Following the APHENITY clinical trial, this extension study looked at the long-term safety and potential benefits of SEPHIENCE. The goal was to see if SEPHIENCE could help people eat more natural protein while keeping their blood Phe levels in a healthy range.



Who took part?

As of the data cutoff date of September 2, 2024, a total of 169 people with PKU joined this study. Participants ranged in age from ~2 months to 55 years old, with a median age of 14. Some people had taken part in the original APHENITY trial, while others were new to SEPHIENCE. This study is still in progress, and the results shown are based on the data collected up to the cutoff date.

How did the study work?

- Everyone took SEPHIENCE once a day for at least one year
- 169 participants were divided into two groups based on their blood Phe levels at the start
 - Those with lower Phe levels ($<360 \mu\text{mol/L}$, $n=102$) were guided through a diet assessment period, where healthcare providers slowly increased the amount of natural protein they could eat, adjusting the diet every two weeks over 26 weeks
 - Those with higher Phe levels ($\geq 360 \mu\text{mol/L}$, $n=49$) continued on treatment without diet changes, but all participants had regular check-ins throughout the study



IMPORTANT SAFETY INFORMATION (cont'd)

How should I take or give SEPHIENCE? (cont'd)

- Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with SEPHIENCE if you have certain side effects.
- Your healthcare provider may change your diet during treatment. Follow your healthcare provider's instructions and do not make any changes to your diet without first talking to your healthcare provider.

Please see full Important Safety Information on [pages 12-13](#), and the [Full Prescribing Information](#).



With SEPHIENCE™, many people were able to eat more natural protein

After 26 weeks of treatment, researchers examined changes in the amount of natural protein participants in the dietary assessment group were able to consume relative to their baseline.

Here's what they found:



97%

increased their dietary Phe intake

With SEPHIENCE, nearly all participants could relax their diet and increase Phe intake during the 26-week assessment period.



70%

met their daily recommended protein needs

Two out of three participants met their age-adjusted natural protein needs while on SEPHIENCE.



~60g

of protein per day

With SEPHIENCE, a 110-lb person could consume approximately 60 grams of natural protein per day after 26 weeks of treatment.

Individual results may vary. This information is based on clinical trial data but is not included in the approved full Prescribing Information for SEPHIENCE. Talk to your healthcare provider if you have any questions.

Study limitations:

This study is “open-label,” meaning both the researchers and participants knew what treatment was being given. Also, participants reported their own protein intake, which may not always be exact. Sometimes diary records were incomplete. Because people of various ages participated and had different starting diets, this may impact our understanding of the results. The APHENITY Extension Study cannot say for sure whether people with PKU can fully relax their diet. However, it shows that SEPHIENCE continues to work over the long term to keep blood Phe levels below 360 $\mu\text{mol/L}$ (6 mg/dL), which for some adults and children with PKU may mean being able to ease dietary restrictions. This is for informational purposes only.

IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE?

SEPHIENCE may cause serious side effects, including:

- **increased bleeding problems.** SEPHIENCE may increase your risk of bleeding. Tell your healthcare provider if you develop any symptoms of bleeding including:
 - bruising or red or purple skin marks
 - menstrual bleeding that is heavier than normal
 - bleeding longer than usual or that does not stop
- **low blood Phe (hypophenylalaninemia).** Low blood Phe levels are common during treatment with SEPHIENCE and can be severe.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).


Sephience™
 sepiapterin
 250 mg | 1000 mg Oral Powder

What to expect with SEPHIENCE

SEPHIENCE has been shown to provide effective management of Phe levels in clinical trials of people with PKU spanning a range of age groups.

SEPHIENCE may cause serious side effects, including an increased risk of bleeding or bruising. One person in the APHENITY trial experienced non-life-threatening bruising and prolonged bleeding, which resolved after treatment was discontinued. Contact your healthcare provider if you notice any unusual bleeding. In children with PKU, it can sometimes cause very low levels of Phe in the blood, so regular monitoring is important. For those taking levodopa, your healthcare provider should monitor for any side effects such as seizures, irritability, or feeling overly stimulated.

Side effects with SEPHIENCE may affect each person differently. Information on the long-term use of SEPHIENCE is limited.



IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)


The most common side effects of SEPHIENCE include: diarrhea, headache, stomach (abdominal) pain, yellow or orange stool (feces), and throat pain.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of SEPHIENCE. Call your healthcare provider for medical advice about side effects.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

Most common side effects



Diarrhea
(7%)




Headache
(7%)



Stomach pain
(5%)



Throat pain
(4%)



Low levels of Phe in the blood—
below the healthy range of
30–120 $\mu\text{mol/L}$ —also called
HypoPhe (4%)



Unusual-colored
feces
(4%)

These are not all the possible side effects of SEPHIENCE™. If you experience any side effects, talk to your healthcare provider.



Talk to your healthcare provider about your treatment and concerns—they're your best guide to managing PKU.

IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)

Before taking or giving SEPHIENCE, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if SEPHIENCE will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if SEPHIENCE passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take SEPHIENCE.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).



Important Safety Information

What is SEPHIENCE?

SEPHIENCE™ (sepiapterin) is a prescription medicine used to lower blood levels of phenylalanine (Phe) in adults and children 1 month of age and older with a certain type of Phenylketonuria (PKU). SEPHIENCE is used along with a Phe-restricted diet. It is not known if SEPHIENCE is safe and effective in children younger than 1 month of age.

What is the most important information I should know about SEPHIENCE?

SEPHIENCE may cause serious side effects, including:

- **increased bleeding problems.** SEPHIENCE may increase your risk of bleeding. Tell your healthcare provider if you develop any symptoms of bleeding including:
 - bruising or red or purple skin marks
 - menstrual bleeding that is heavier than normal
 - bleeding longer than usual or that does not stop
- **low blood Phe (hypophenylalaninemia).** Low blood Phe levels are common during treatment with SEPHIENCE and can be severe.

The most common side effects of SEPHIENCE include: diarrhea, headache, stomach (abdominal) pain, yellow or orange stool (feces), and throat pain.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of SEPHIENCE. Call your healthcare provider for medical advice about side effects.

Before taking or giving SEPHIENCE, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if SEPHIENCE will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if SEPHIENCE passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take SEPHIENCE.

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. SEPHIENCE and other medicines may affect each other causing side effects. SEPHIENCE may affect the way other medicines work, and other medicines may affect how SEPHIENCE works. Especially tell your healthcare provider if you take:

- a medicine that contains levodopa
- an antifolate medicine
- another medicine also used to lower Phe
- medicines used to treat erectile dysfunction (phosphodiesterase 5 inhibitors or PDE5)

Know the medicines you take. Keep a list of them with you to show your healthcare provider and pharmacist when you get a new medicine.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).



How should I take or give SEPHIENCE?

- Take or give SEPHIENCE exactly as your healthcare provider tells you. Your healthcare provider will tell you how much SEPHIENCE to take or give.
- Your healthcare provider may change your or your child's dose of SEPHIENCE depending on your weight and age.
- Take or give SEPHIENCE 1 time each day with food. You must take or give more food after the SEPHIENCE dose.
- SEPHIENCE comes as a packet containing oral powder.
 - For SEPHIENCE doses **less than** 1,000 mg: mix SEPHIENCE oral powder with water or apple juice before giving or taking.
 - For SEPHIENCE doses 1,000 mg or **greater**: SEPHIENCE oral powder can be mixed with water, apple juice, strawberry jam, or applesauce before taking or giving.
 - **See the detailed Instructions for Use that comes with SEPHIENCE for information about the correct way to prepare, take, or give a dose of SEPHIENCE oral powder.**
- If you or your child miss a dose of SEPHIENCE, take or give it as soon as possible. Do not take 2 doses in a day. Continue with your normal dosing schedule the following day.
- During treatment with SEPHIENCE, you or your child will have regular check-ups with your healthcare provider.
- Your healthcare provider will do blood tests before and during your treatment with SEPHIENCE to check your blood Phe levels.
- Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with SEPHIENCE if you have certain side effects.
- Your healthcare provider may change your diet during treatment. Follow your healthcare provider's instructions and do not make any changes to your diet without first talking to your healthcare provider.

To report an adverse event, please call [1-866-562-4620](tel:1-866-562-4620) or email at usmedinfo@ptcbio.com. You may also report side effects to FDA at [1-800-FDA-1088](tel:1-800-FDA-1088) or at www.fda.gov/medwatch.

Please see [Full Prescribing Information](#).

Once-daily dosing

Taking SEPHIENCE

- Take SEPHIENCE once a day, with food
 - Additional food intake is required after dosing
- It's best to take SEPHIENCE at the same time each day
- Your healthcare provider will tell you the right dose based on your (or your loved one's) body weight and age

Mixing SEPHIENCE

- Comes in **250 mg** and **1000 mg** powder packets
- Mix with water, apple juice, or soft food (only for doses ≥ 1000 mg) like applesauce or strawberry jam
- May cause staining when in contact with certain surfaces. To minimize this risk, use as directed

Personalized dose tracking

Use the SEPHIENCE dosing calculator on our website (**for informational purposes only**). **This tool does not replace medical advice.** Always follow your healthcare provider's instructions.

**Try the
dosing calculator**

IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. SEPHIENCE and other medicines may affect each other causing side effects. SEPHIENCE may affect the way other medicines work, and other medicines may affect how SEPHIENCE works. Especially tell your healthcare provider if you take:

- a medicine that contains levodopa
- an antifolate medicine
- another medicine also used to lower Phe
- medicines used to treat erectile dysfunction (phosphodiesterase 5 inhibitors or PDE5)

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).



Record the dose as discussed with your healthcare provider and update it as needed

If the daily dose is <1000 mg

DAILY DOSE mg

of 250 mg packets:

of 1000 mg packets:

OR

Mix in

ml

of water or apple juice

PRESCRIBED VOLUME

ml

to be administered

If the daily dose is ≥1000 mg

DAILY DOSE mg

of 250 mg packets:

of 1000 mg packets:

AND
OR

Mix in

ml
or
Tbsp

of water,
apple juice,
strawberry jam,
or applesauce

ADMINISTER THE FULL VOLUME



See the next page for step-by-step instructions on how to take SEPHIENCE.

IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about SEPHIENCE? (cont'd)

Know the medicines you take. Keep a list of them with you to show your healthcare provider and pharmacist when you get a new medicine.

How should I take or give SEPHIENCE?

- Take or give SEPHIENCE exactly as your healthcare provider tells you. Your healthcare provider will tell you how much SEPHIENCE to take or give.
- Your healthcare provider may change your or your child's dose of SEPHIENCE depending on your weight and age.
- Take or give SEPHIENCE 1 time each day with food. You must take or give more food after the SEPHIENCE dose.

Please see full Important Safety Information on [pages 12-13](#), and the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder

Follow these steps to take SEPHIENCE



Prepare

Mix the correct number of packets with water, apple juice, or soft food (only for doses ≥ 1000 mg), such as applesauce or strawberry jam. These mixing options were studied in clinical trials. For additional mixing options, please check with your healthcare provider.



Mix until combined

- Stir liquids for at least 30 seconds
- Stir soft foods for at least 60 seconds

Note: Oral powder is not expected to dissolve completely. This is normal.



Use as directed

- Drink or swallow the full amount
- For doses < 1000 mg: Use an oral syringe to draw up the prescribed dose and administer



Rinse & finish

- Rinse the cup or syringe with water or juice
- Swallow the rinse until clean to make sure you get the full dose



If you need to store it

- If not taken right away, cover and store the mix:
 - In the refrigerator (36°F to 46°F / 2°C to 8°C) for up to 24 hours
 - At room temperature (68°F to 77°F / 20°C to 25°C) for up to 6 hours
- Stir again before using (see step 2) for at least 30 to 60 seconds

Refer to Prescribing Information (page 18) for [Instructions for Use](#).

IMPORTANT SAFETY INFORMATION (cont'd)

How should I take or give SEPHIENCE? (cont'd)

- SEPHIENCE comes as a packet containing oral powder.
 - For SEPHIENCE doses **less than** 1,000 mg: mix SEPHIENCE oral powder with water or apple juice before giving or taking.

Please see full Important Safety Information on [pages 12–13](#), and the [Full Prescribing Information](#).

Please see full Important Safety Information on [pages 12-13](#), and the [Full Prescribing Information](#).





WE'VE GOT YOU COVERED

Not actual patients.

Enroll in PTC Cares to get connected with your dedicated Case Manager.

PTC Cares Case Managers guide families through treatment, insurance, and medication access. Think of your Case Manager as your personal guide!

Have questions?

Contact PTC Cares at
1-844-4PTC-CARES
(1-844-478-2227)

Monday to Friday, 8am to 6pm (EST)

My Case Manager

Name

Phone Number



How we can help with PKU treatment costs

Our Case Managers explore various options to help access the PKU medication your healthcare provider prescribes. PTC Therapeutics offers several programs to help those enrolled in PTC Cares™.



The Copay Assistance Program*†

May help qualified families who have commercial insurance get help covering out-of-pocket costs associated with treatment. Patients who qualify may pay as little as \$0 copay.



The Bridge Program*

May provide patients currently on a PTC medication with a temporary supply free of charge while they await verification of insurance coverage.



Patient Assistance Program (PAP)

May help patients without insurance, who are eligible, get access to treatment.

*Low to no out-of-pocket costs for most patients who qualify.

†Not valid for prescriptions eligible to be reimbursed, in whole or in part, by Medicaid or Medicare (including Medicare Part D), or where prohibited by law.

Join PTC Cares, treatment support for people and families living with PKU

To get started, simply access and sign the PKU Patient Consent for PTC Cares Program online at PTCCares.com.



A PKU TREATMENT THAT HELPS FIND

LESS LIMITS. MORE POSSIBILITIES.

See what SEPHIENCE can offer.



Flexible once-daily dosing



Shown to reduce Phe levels
across all ages, from children
to adults



Support and resources
available at every step



Proven safety profile

Visit [SEPHIENCE.com](https://sephience.com) or speak
with your healthcare provider for
more information

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about
SEPHIENCE?

SEPHIENCE may cause serious side effects, including:

- **increased bleeding problems.** SEPHIENCE may increase your risk of bleeding. Tell your healthcare provider if you develop any symptoms of bleeding including:
 - bruising or red or purple skin marks
 - menstrual bleeding that is heavier than normal
 - bleeding longer than usual or that does not stop
- **low blood Phe (hypophenylalaninemia).** Low blood Phe levels are common during treatment with SEPHIENCE and can be severe.

Please see full Important Safety
Information on [pages 12-13](#), and
the [Full Prescribing Information](#).


Sephience™
sepiapterin
250 mg | 1000 mg Oral Powder



PTC Cares, PTC Cares Logo and SEPHIENCE are
trademarks of PTC Therapeutics, Inc. or affiliates.
© 2025 PTC Therapeutics, Inc. All Rights Reserved.
US-SEP-0223 10/25